

# Applied DC Magnetic Fields Cause Alterations in the Time of Cell Divisions and Developmental Abnormalities in Early Sea Urchin Embryos

Michael Levin\* and Susan G. Ernst

Department of Biology, Tufts University, Medford, Massachusetts

Most work on magnetic field effects focuses on AC fields. The present study demonstrates that exposure to medium-strength (10 mT–0.1 T) static magnetic fields can alter the early embryonic development of two species of sea urchin embryos. Batches of fertilized eggs from two species of urchin were exposed to fields produced by permanent magnets. Samples of the continuous cultures were scored for the timing of the first two cell divisions, time of hatching, and incidence of exogastrulation. It was found that static fields delay the onset of mitosis in both species by an amount dependent on the exposure timing relative to fertilization. The exposure time that caused the maximum effect differed between the two species. Thirty millitesla fields, but not 15 mT fields, caused an eightfold increase in the incidence of exogastrulation in *Lytechinus pictus*, whereas neither of these fields produced exogastrulation in *Strongylocentrotus purpuratus*. *Bioelectromagnetics* 18:255–263, 1997. © 1997 Wiley-Liss, Inc.

**Key words:** sea urchin; static magnetic field; gastrulation; development; mitotic cycle; teratogenic effects

## INTRODUCTION

The resurgence of interest in the interactions between electromagnetic fields and biological systems has focused mainly on AC (time-varying) fields. However, there have been studies showing that DC (static) magnetic fields can also interact with living systems at various levels. Effects on in vitro biochemical reactions have been reported [Kim, 1976; Adamkiewicz and Pilon, 1983; Adamkiewicz et al., 1987; EPA, 1990; Markov et al., 1992; Richardson et al., 1992; Harkins and Grissom, 1994]. Perhaps most interestingly, it has been shown that a 50 mT DC magnetic field can alter the structure of poly-L-lysine [Verma and Goldner, 1996].

Behavioral effects of DC fields have also been noted. For example, strong static magnetic fields are avoided by mice and worker ants [Kermarrec, 1981], although a DC magnetic field of about 0.1 mT apparently increases bee life span by more than 60% [Martin et al., 1989]. Weak static fields affect the choice of motion of flatworms, snails, and paramecia [Dubrov, 1978; Martin et al., 1989], medium-strength static fields have been used as conditioning stimuli for bees and rabbits [Kholodov, 1971; Walker and Bitterman, 1989], and a 7 mT static field disrupted honeybee danc-

ing [Tomlinson et al., 1981]. Medium-strength DC fields act as a general stressor in mice [Laforge et al., 1978, 1986].

Physiological effects, such as changes in leukocyte count in mice [Barnothy, 1957], disruption of the mammalian menstrual cycle [Kholodov, 1973], reduced respiration in cultured embryonic and sarcoma cells [Pereira et al., 1967], alterations in growth rate of plants and bacteria [Dycus and Shultz, 1964; Pittman, 1972; Singh et al. 1994], and changes in aging rates [Kholodov, 1971; Bellossi, 1986], have also been reported. Likewise, morphological and histochemical alterations in rat spermatogenesis [FBIS, 1983] and central nervous system (CNS) microstructure [Abdullakhodzhayeva and Razykov, 1986], cytological changes

---

Contract Grant sponsor: Howard Hughes Medical Institute, Initiative in Undergraduate Education; Contract Grant sponsor: U.S. EPA; Contract Grant number: CR820-301.

Michael Levin is now at the Department of Genetics, Harvard Medical School.

\*Correspondence to: Michael Levin, Department of Genetics, Harvard Medical School, Boston, MA 02115.

Received for review 9 August 1995; Revision received 12 July 1996

in paramecia [Kogan et al., 1967], reduction of X irradiation-induced mortality [Barnothy, 1963], retardation of wound healing [Beischer, 1964], and abnormal mitotic figures and nuclei [Linskens and Smeets, 1978; Mastryukova and Rudneva, 1978; FBIS, 1983] appear to be caused by exposure to medium-strength to high-strength static magnetic fields.

Some attention has been focused on interactions between static fields and processes involved in carcinogenesis and tumor formation. As is true for the effects of AC fields [reviewed in Bates, 1991], the effects of static fields on tumors can appear contradictory, depending on field parameters. Fields of 730 mT cause cell degeneration in several types of tumor cells [Kim, 1976]. Gross [1962] found that a 400 mT field increased the rate of death from transplanted tumors in mice, yet treatment of H2712 mouse tumor cells with a 3.8 T field (with a 1.2 T/mm gradient) caused significant inhibition of the ability to infect a healthy host [Weber and Cerilli, 1971]. Recent studies have shown that the oncogene *c-fos* can be induced by 0.2 T static field in cultured mammalian cells [Hiraoka et al., 1992]. DC magnetic fields (0.73 T) applied to tumor cell suspensions can cause a sharp reduction in cell number [König et al., 1981].

Especially interesting are the reports that static fields are able to alter embryonic development and morphogenesis, because, in addition to the basic question of mechanisms of field-biosystem interaction, these effects provide an opportunity to learn more about developmental mechanisms. It has been reported that 1 T fields are lethal to young mice [Kholodov, 1971] and that a 14 T field stopped sea urchin development but did not affect *Drosophila* and mouse development [Kholodov, 1971]. A weaker, 420 mT field, caused embryo death and dissolution in the wombs of mice [Kholodov, 1971]. Klueber [1981] showed that 5 mT static magnetic fields produced dramatic teratogenic effects in the eye and nervous system of developing chick embryos. Static magnetic fields of 0.4 mT retarded development of the pigeon embryo, and exposure of chick embryos to a 500 mT field for just 1 h produced poor brain development with an open neural tube, shortening of the embryonic long axis, and slight heart displacement [Joshi et al., 1978]. Neurath [1968, 1969] showed that, in many organisms, gastrulation is halted in fields with a gradient of 8.35 T/cm, and *Drosophila* cuticular abnormalities resulted from brief exposures to static magnetic fields [Ho et al., 1992]. One tesla T DC fields caused axial anomalies in frog embryos [Ueno, 1984]. Regeneration, a nonembryonic example of large-scale morphogenesis, is also affected by DC fields, which accelerate tail regeneration in tadpoles; the effect exhibits exposure time and field

strength dependence [Kudokozev and Baranovskiy, 1988].

We performed a series of experiments to examine the effects of medium-strength static magnetic fields on the development of sea urchin embryos. The sea urchin is an excellent and well-studied developmental system, and the presence of static magnetic field effects on its development would afford a tractable model for studying field-cell interactions, as well as the normal processes of development, by providing a new perturbing factor. In one sense, static field effects are more interesting, because, unlike AC fields, they cannot cause ionic currents. Although our initial studies [Levin and Ernst, 1995] showed that weak AC magnetic fields can affect the mitotic timing of sea urchin embryos and Kholodov [1971] reported that a high-strength (14 T) static magnetic field arrests sea urchin development, there have been very few experimental studies of applied medium-strength static field effects on sea urchin embryogenesis. In the present paper, we report that such fields are able to cause a delay in the mitotic cycle of early embryos and to increase greatly the incidence of exogastrulation, a well-characterized developmental abnormality in sea urchins. Thus, we show that, in the sea urchin model, static magnetic fields are a potent teratogen and that the mitotic cycle is sensitive to these low-energy fields as well as to AC fields.

## MATERIALS AND METHODS

### Animals, Gametes, and Embryos

*Strongylocentrotus purpuratus* and *Lytechinus pictus* were purchased from Marinus, Inc. (Long Beach, CA). Animals were maintained in aquaria at 9 °C and were induced to spawn by intracoelomic injection of 0.5 M KCl. Semen was collected dry from the genital pores with a Pasteur pipette and held undiluted in a tube on ice until fertilization. Eggs were collected by inverting the spawning females onto beakers of Millipore-filtered sea water (MPFSW). The suspension of eggs was filtered through several layers of cheesecloth and settled on ice through fresh MPFSW three times. Eggs were suspended to a final concentration of 1–2% (V:V) in MPFSW.

Fertilization was accomplished by adding a freshly prepared dilute sperm suspension to the eggs to result in a final sperm concentration of about 1:10 000 (V:V). Successful fertilization was determined by elevation of the fertilization membrane, generally within 90 s of sperm addition. Fertilization was greater than 95% in all experiments. Experiments were carried out with eggs produced from several females to minimize effects of individual differences. Embryos in 250 ml




Fig. 1. Apparatus for magnetic field exposure. The culture is exposed to a static field produced by two attracting ceramic magnets. Asterisks represent embryos or eggs.

beakers were cultured with stirring in an incubator at 13–16 °C, depending on the species.

**Experimental Design for Static Field Exposure**

The static magnetic fields were produced by a parallel pair of attracting rectangular ceramic magnets positioned opposite each other, with the sample of sperm, eggs, or embryos between them (Fig. 1). Field strength was measured with a Gaussmeter (Walker Magnetics; model MG-4D) at the midpoint of the culture. Field strength at the outer edges of the culture was no more that ±15% of this value.

Styrofoam blocks were used to insulate the stirring motors from the cultures, to guard against possible differential heating effects FROM the motors. Temperature between the test and control cultures in individual experiments did not differ by more than ±0.5 °C, as determined by continuous monitoring. The magnetic environment of the incubator was investigated with the Gaussmeter and was determined not to be different from the ambient geomagnetic field, within ±0.01 mT. The stirring motors produced no detectable stray fields at the location of the cultures (35 cm away from motors). All supporting material within the incubator was made of nonferrous material to prevent unwanted leakage of the fields toward the control culture. All aspects of culture, except for the presence of magnets, were the same in the exposed and control samples (including beakers, stirring motors, etc.).

**Sampling and Data Collection**

During the continuous experiments, samples of about 200 embryos were taken without interruption of

**TABLE 1. *L. pictus* Embryos Were Exposed to a 30 mT Static Field Immediately After Fertilization. Twenty-six Hours Later, Samples from the Control and Exposed Batches Were Scored for Hatching (Absence of Fertilization Membrane)**

	Control	Exposed
Hatched	82%	36%
Not hatched	15%	60%
Arrested before hatching	3%	4%
Total embryos	109	111

field exposure approximately every 15 min, fixed in 3% formaldehyde, and scored for the number of blastomeres, the presence of the fertilization membrane, or the position of the gut with the aid of a Nikon microscope.

**RESULTS**

**Exposure to Static Fields Delays Hatching**

The first series of experiments was designed to maximize the possibility of detecting effects arising from exposure to static magnetic fields. Immediately following fertilization, *L. pictus* embryos were divided into two equal volumes; one culture was exposed to a 30 mT static field (Fig. 1) for the duration of the experiment, and the other culture received no exposure aside from the ambient geomagnetic field. All other conditions of culture remained the same. At 26 h postfertilization, samples of each culture were taken and scored for percentage of embryos hatched. Hatching is an easily recognizable developmental event resulting from blastula-stage embryos acquiring motility and secreting an enzyme that digests the fertilization membrane [reviewed in Okazaki, 1975]. The results are summarized in Table 1 and demonstrate that, at 26 h, 82% of the control embryos had hatched, whereas only 36% of the exposed embryos had done so ( $P < .01$ ). Additional experiments with *L. pictus* and *S. purpuratus* revealed that, in both species, exposure to 30 mT static magnetic fields significantly delays hatching relative to control groups (data not shown).

**Exposure to Static Field Delays the First and Second Cell Divisions of *S. purpuratus***

Having seen that exposure to the field is able to delay hatching time, we hypothesized that this was due to an increase in the length of the cell cycle rather than to a delay of the hatching mechanism itself. Consequently, we studied the effect of exposure on the time of the first two cell divisions. A batch of eggs was split in two immediately following fertilization, and one culture was exposed to a 30 mT magnetic field (Fig. 1).

**TABLE 2. *L. pictus* Embryos Were Exposed to a 30 mT Static Field Immediately After Fertilization. At 3.5 Hours Post-fertilization, Samples from the Control and Exposed Cultures Were Scored for the Number of Blastomeres**

	Control	Exposed
1 cell	9%	36%
2 cell	91%	64%
4 cell	0%	0%
Other	0%	0%
Total embryos	104	111

At 3.75 h, approximately 100 embryos from each culture were scored for cell division. Four times as many exposed eggs relative to controls remained undivided (Table 2). These results demonstrate that the field increases the length of time between completed cell divisions.

To gain more information on which phases of the cell cycles were affected and to have a more quantitative way of determining how the magnitude of the delay varied with field parameters, the continuous cultures were sampled and scored periodically, thereby recording how many cell divisions had taken place in a representative sample of the embryos. This made it possible to plot the number of divisions as a function of time and then to compare exposed and control batches of embryos. Cell divisions in early sea urchin embryos are well synchronized; we have previously demonstrated that when a batch of fertilized eggs is split and sampled every 15 min, the maximum endogenous variation of the time of cell division is 3 min [Levin and Ernst, 1995]. Thus, differences greater than  $\pm 3$  min were taken to be significant in the experiments discussed below.

Because the endogenous variation is known, it was possible to study the effects of exposure to the 30 mT static field on the time of the first two cell divisions. After fertilization, the embryos were split into two cultures; one was immediately exposed to the field throughout the experiment, and the other served as a control. The cultures were sampled every 15 min without interruption of the field, and roughly 200 embryos from each culture were scored for the number of blastomeres. The results are shown in Figure 2. By calculation of the time difference between the midpoints of each cell division phase on the plot, it was shown that the field induces an insignificant delay (1 min) in the first cell division and a small but significant delay of 6 min in the second.

#### Effects of Static Field Applied to Sperm on Timing of First/Second Cell Divisions

Having seen that exposure to the field resulted in a delay in the time of cell division, we tested the possi-




Fig. 2. Exposure of embryos to a 30 mT field immediately at fertilization. Solid line and asterisks indicate the curve for the control culture, and dashed line and open circles indicate the curve for the exposed culture. The field produces an insignificant delay (1 min) in the first cell division and a small but significant (6 min) delay in the second relative to controls.

bility that some mechanism within the sperm is sensitive to the field. A 30 mT static field was applied to the undiluted sperm sample of *S. purpuratus* for 1 h. The field was removed immediately prior to dilution of the sperm sample for fertilization. The resulting culture and a control were incubated without exposure to any field, and samples were taken, scored, and plotted as discussed above. No significant effects on the time of the first two cell divisions were seen (Fig. 3). There was also no obvious decrease in the ability of sperm to fertilize eggs; greater than 95% fertilization was achieved within 5 min, the same as in the control sperm samples. The same result was observed for *L. pictus* (data not shown). However, because sperm were not limiting, we cannot rule out the possibility that a fraction of the sperm was affected by exposure to the field.

#### Magnitude of Cell Division Delay Is a Function of Timing of Field Exposure

Because exposure of sperm to a 30 mT static field had no measurable effect, we were interested to see whether prefertilization eggs were sensitive to the field or the field only affected activated eggs or developing embryos. It was also possible to study the relationship between the time of onset of field exposure and the magnitude of the delay. A 30 mT static field was applied to the experimental cultures of *S. purpuratus* starting at various times relative to fertilization and lasting throughout the experiment. The cell division time profiles were calculated as discussed above.




Fig. 3. Exposure of sperm alone to a 30 mT static field for 1 h. Solid line and asterisks indicate the curve for the control culture, and dashed line and open circles indicate the curve for the exposed culture. The field has no significant effect on the duration of the first two cell divisions.




Fig. 4. Exposure of embryos to a 30 mT field at 15 min prefertilization. Solid line and asterisks indicate the curve for the control culture, and dashed line and open circles indicate the curve for the exposed culture. The exposure results in a 17 min delay in each of the first two cell divisions relative to controls.

When eggs were exposed to the field beginning at 45 min prefertilization, the first cell division was delayed 4 min and in the exposed cultures the second division was 6 min behind that measured for the control culture (although these figures are larger than the 3 min endogenous difference between control cultures, the limited number of experiments precludes determining whether these differences are significant by a comprehensive statistical analysis). Exposure that began at 30 min before fertilization caused a 10 min delay in the first cell division and a 13 min delay in the second, whereas an exposure begun at 15 min prefertilization caused a 17 min delay in the timing of both cell divisions (Fig. 4). When eggs were exposed starting at 6 min prefertilization, there was a 14 min delay in the first cell division and a 15 min delay in the second division. In cultures exposed immediately following fertilization, there was an insignificant delay for the first cell division and a 6 min delay for the second cell division. These results are summarized in Figure 5. When this series of experiments was repeated with *L. pictus*, the same type of relationship was observed between timing of exposure and magnitude of delay, except that the optimal time of exposure was shown to be 30 min prefertilization and the delay obtained at that exposure was 22 min (data not shown).

**Morphological Effects**

Given the reported teratological effects of magnetic fields (reviewed above), we were interested in determining whether our field exposures had any such effects. For *L. pictus* cultures exposed to a 30 mT field




Fig. 5. A nonlinear relationship between exposure timing and magnitude of cell cycle delay was observed. The optimal exposure time is 15 min prefertilization, which results in a delay of 17 min in each cell division.

continuously from fertilization and incubated for 48–94 h, we noticed an apparent increase in the incidence of exogastrulation. Exogastrulation is a well-known developmental abnormality in sea urchins in which the archenteron, the primitive gut, evaginates, forming outside the embryo, instead of invaginating [Nocente-

**TABLE 3. *L. pictus* Embryos Were Exposed to a 30 mT Static Field Immediately After Fertilization. The Exposure Lasted Throughout Development. Two to Three Days Later, Samples of the Control and Exposed Cultures Were Scored for the Incidence of Exogastrulation**

	Control	Exposed
Normal gastrula	98%	88%
Exogastrula	2%	12%
Total embryos	109	111




McGrath et al., 1991]. To test directly the effect of a 30 mT static field on *L. pictus* gastrulation, embryos were fertilized and the culture was split in half, with one half serving as a control and the other half exposed to the field. Gastrulation is normally initiated about 24 h postfertilization in *L. pictus*, followed by the morphological differentiation of the gut. Cultures were scored for exogastrulation at 2–3 days.

In exposed cultures, the incidence of exogastrulation rose from control values of 1–2% to as high as 16%. In seven experiments, the increase in exogastrulation was between twofold and eightfold, on average sixfold, above that of controls. The results of one such experiment ( $\chi^2 = 8.94$ ,  $P < .05$ ) are shown in Table 3, and examples of exogastrulated embryos produced by exposure to a 30 mT static field are shown in Figure 6. A 0.39 mT 60 Hz AC field also resulted in a threefold increase in exogastrulation, whereas 0.195 and 0.016 mT AC fields did not measurably alter the incidence of exogastrulation (data not shown). For *S. purpuratus* embryos exposed to the same fields, we found no increase in the incidence of exogastrulation.

Interestingly, a morphological abnormality not reported before in either species and never observed in controls was found in the eggs exposed to a 30 mT static field. This abnormality, shown in Figure 7, consisted of embryonal collapse along one axis, resulting in a flat disk rather than the normal sphere. In three experiments in which fertilized eggs were exposed to a 30 mT static field at 45 min, 30 min, or 0 min pre-fertilization, and continuing throughout the subsequent 48–94 hours, approximately 1% of the eggs collapsed.

## DISCUSSION

In this study we observed that a 30 mT static magnetic field applied to sea urchin eggs produced alterations in the time of cell division and induced two developmental abnormalities, exogastrulation and collapsed embryos. These results are surprising, and at present we do not have a unifying model for a mechanism that could account for the diverse effects.



**Fig. 6.** Field-induced exogastrulated (arrowheads) embryos. *L. pictus* embryos were exposed to a 30 mT static magnetic field at fertilization, and the field was maintained throughout the experiment. **A** and **B** show examples of field-induced exogastrulated embryos. **C** is a normal, nonexposed, embryo at the same age.

Exposure to a 30 mT static field resulted in delays in the time of the first two cell divisions of sea urchin embryos. Because the slopes of the division profile curves are roughly equal between the exposed and control embryos (e.g., see Fig. 4), it can be concluded that the field increases the time spent in the G1, G2, or S phases of the cell cycle, rather than slowing down cytokinesis itself. The delay in the time of cell division




Fig. 7. Field-induced collapsed embryos. *L. pictus* embryos were exposed to a 30 mT static magnetic field at fertilization, and the field was maintained throughout the experiment. **A** shows an example of field-induced collapsed embryos. **B** is a normal, nonexposed, embryo at the same age.

seems to be an effect of the static field on the egg and not on the sperm.

It is interesting to note that the static field produces relatively equal delays for the first and second cell divisions (Fig. 5), compared to our earlier studies, in which exposure to an AC field produced much larger accelerations for the second cell division than for the first [Levin and Ernst, 1995]. The AC field seems to shorten each mitotic cycle, whereas the static field appears to act once, before the first cell division and, most likely, before fertilization, because earlier preferential exposures have greater effects.

The static field effect does not show a simple dose-effect relationship with cell division time, because longer exposures do not necessarily produce a more pronounced effect than shorter exposures. Rather, a bell-curve relationship around a maximal value is observed, which may represent some sort of habituation process within the egg. This observation is potentially quite important, because most EMF-exposure guidelines are designed to limit the field magnitude and the amount of time a person spends within a field. Our data suggest that the crucial parameter may be not the length of a biological system's exposure to the field (because, as is shown in Fig. 5, shorter exposures can produce a larger effect) but, rather, the relationship between timing of exposure and key biological events. The exposure timing that produces the maximal delay in *S. purpuratus* is 15 min prefertilization. The 17 min delay may be the maximum effect that can be achieved by this field, because it is the same for both divisions, whereas the other experiments show slightly greater delays for the second division. The same bell-shaped relationship was seen for *L. pictus* (data not shown), resulting in a slightly different optimal timing (30 min prefertilization) and a somewhat greater delay (22 min) than for *S. purpuratus*.

*L. pictus* embryos exposed to static and AC fields exhibited up to an eightfold increase in the incidence of exogastrulation, whereas none of the applied fields tested had this effect on *S. purpuratus* embryos. This is consistent with the fact that the natural incidence of exogastrulation in *S. purpuratus* is much lower than in *L. pictus* and with the fact that *L. pictus* exogastrulates at a lower LiCl concentration than does *S. purpuratus* [Nocente-McGrath et al., 1991]. Thus, this differential sensitivity to magnetic fields likely reflects a genetic difference between the two species.

The collapsed embryos represent an unknown phenomenon. This was observed in single-cell eggs as well as in cultures of hatching embryos. It is unknown, however, whether the collapsed embryos seen in mid-blastula cultures represent embryos that collapsed before the first cell division or embryos that collapsed at some later time. This defect has been observed in unfixed embryos, ruling out artifacts caused by fixation. No other known teratogenic agent is known to produce similar effects.

The effects described were most likely due to the applied DC field. The small stirring motors produced no detectable AC fields at the level of the magnets. Though we cannot formally completely rule out the possibility of very weak eddy currents being induced in the magnets by stray AC fields, this is extremely unlikely to have caused the effects described above; previous experiments have indicated that somewhat

weaker ceramic magnets used in exactly the same experimental setup have no effects (data not shown). The static field cannot induce heating and is several orders of magnitude weaker than fields that have been shown to cause changes in membrane electrical properties [Kholodov, 1971]. Furthermore, ferrous molecules are not known to occur in sea urchins. However, our results demonstrate that static fields containing very little energy (of the same order of magnitude as average kinetic energy due to thermal motion) can significantly affect important biological processes. DC magnetic fields can, however, alter the velocity of motion of ions.

Under appropriate conditions, small changes in the behavior of ions could cause significant effects. For example, alterations in velocity would affect the interactions between ions and receptor channels. Similarly, small changes in the behavior of ions that are components of signal transduction pathways could produce dramatic changes. Interestingly, lithium, the classic inducer of exogastrulation in sea urchins [reviewed in Nocente-McGrath et al., 1991], is proposed to act by secondary-messenger pathways in several different cell types [reviewed in Berridge et al., 1989]. Still, there is no immediately obvious mechanism that can account for the various developmental effects we have observed. Some possibilities are induced changes in trajectories of moving ions such as  $\text{Ca}^{2+}$  near membranes, electric currents induced by the motion of the conductive cytoplasm of the urchins through the static field as they are being stirred, and conformational changes in the structure of regulatory proteins [Verma and Goldner, 1996].

## ACKNOWLEDGMENTS

This work was supported by the Howard Hughes Medical Institute, Initiative in Undergraduate Education and the U.S. EPA, under assistance agreement CR820-301 to the Center for Environmental Management, Tufts University. These studies do not necessarily reflect the views of the EPA and no official endorsement should be inferred. We thank Jorge Gonzalez for helpful discussions and Drs. Verma and Goldner for sharing their unpublished results with us.

## REFERENCES

- Abdullakhodzhayeva MS, Razykov SP (1986): Structural changes in the central nervous system caused by exposure to permanent magnetic field. *Bull Exp Biol Med* 11:600–602.
- Adamkiewicz VW, Pilon DR (1983): Magnetic stimulation of polysaccharide accumulation in cultures of *Streptococcus mutans*, *Can J Microbiol* 29:464–467.
- Adamkiewicz VW, Bassous C, Morency D, et al. (1987): Magnetic response in cultures of *Streptococcus mutans*. *Exp Biol* 46:127–132.
- Barnothy MF (1957): Influence of magnetic fields upon the leukocytes of the mouse. In Quastler H, Morowitz HJ (eds): "Proceedings of the First National Biophysics Conference." New Haven, CT: Yale University Press.
- Barnothy MF (1963): Reduction of radiation mortality through magnetic pretreatment. *Nature* 200:279–280.
- Barnothy MF (ed) (1967): "Third International Biomagnetic Symposium." Chicago: Chicago University Press.
- Bates MN (1991): Extremely low frequency electromagnetic fields and cancer: The epidemiologic evidence. *Environ Health Perspect* 95:147–156.
- Beischer DE (1964): Biomagnetics. *Ann NY Acad Sci* 134:454–458.
- Bellossi A (1986): Effect of static magnetic fields on survival of leukemia-prone AKR mice. *Radiat Environ Biophys* 25:75–80.
- Berridge MJ, Downes CP, Hanley MR (1989): Neural and developmental actions of lithium: A unifying hypothesis. *Cell* 59:411–419.
- Dubrov AP (1978): "The Geomagnetic Field and Life." New York: Plenum Press.
- Dycus A, Schultz A (1964): A survey of the effects of magnetic environments on seed germination and early growth. *Plant Physiol* 35:29.
- EPA (1990): "Evaluation of the Potential Carcinogenicity of Electromagnetic Fields." Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development.
- FBIS (Foreign Broadcast Information Service) (1983): USSR report (life sciences): Effects of nonionizing electromagnetic radiation. No. 2, JPRS 84221, National Technical Information Service. Washington, DC: U.S. Department of Commerce.
- Gross L (1962): Effect of magnetic field on tumor immune responses in mice. *Nature* 195:662–663.
- Harkins TT, Grissom CB (1994): Magnetic field effects on  $\text{B}_{12}$  ethanolamine ammonia lyase: Evidence for a radical mechanism. *Science* 263:958–959.
- Hiraoka M, Miyakoshi J, Li YP, et al. (1992): Induction of c-fos gene expression by exposure to a static magnetic field in HeLaS3 cells. *Cancer Res* 52:6522–6523.
- Ho M-W, Stone TA, Jerman I, Bolton J, Bolton H, Goodwin BC, Saunders PT, Robertson F (1992): Brief exposures to weak static magnetic fields during early embryogenesis cause cuticular pattern abnormalities in *Drosophila* larvae. *Phys Med Biol* 37:1171–1179.
- Joshi MV, Khan MZ, Damle PS (1978): Effect of magnetic field on chick morphogenesis. *Differentiation* 10:39–43.
- Kermarrec A (1981): Sensitivity to artificial magnetic fields and avoiding reaction. *Insect Soc* 28:40–46.
- Kholodov YuA (1971): "Influence of Magnetic Fields on Biological Objects." JPRS 63038, National Technical Information Service. Washington, DC: U.S. Department of Commerce.
- Kholodov YA (1973): "Magnetism in Biology." JPRS 60737, National Technical Information Service. Washington, DC: U.S. Department of Commerce.
- Kim YS (1976): Some possible effects of static magnetic fields on cancer. *Tower Int Technomed Inst Life Sci* 5:11–28.
- Klueber KM (1981): The teratogenicity of low-level magnetic fields in the developing chick embryo. *Anat Rec* 199:144a.
- Kogan AB, Dorojkina LI, Ostapenko LV (1967): Cytological changes of *Paramecium caudatum* in a magnetic field. In Barnothy MF (ed): "Third International Biomagnetic Symposium." Chicago: Chicago University Press.
- König HL, Krueger AP, Lang S, Sönning W (1981): "Biologic Effects of Environmental Electromagnetism." New York: Springer-Verlag.

- Kudokozev VP, Baranovskiy AE (1988): Influence of constant magnetic fields on the regeneration of the tail in amphibian larvae. *Ontogenez* 5:508–512.
- Laforge H, Moisan M, Champagne F, Seguin M (1978): General adaptation syndrome and magnetostatic field: Effects on sleep and delayed reinforcement of low rate. *J Psychol* 98:49–55.
- Laforge H, Sadeghi MR, Seguin MK (1986): Magnetostatic field effect: Stress syndrome pattern and functional relation with intensity. *J Psychol* 120:299–304.
- Levin M, Ernst SG (1995): Applied AC and DC magnetic fields cause alterations in the mitotic cycle of early sea urchin embryos. *Bioelectromagnetics* 16:231–240.
- Linskens HF, Smeets P (1978): Influence of high magnetic fields on meiosis. *Experientia* 34:42.
- Markov MS, Wang S, Pilla AA (1992): Weak pulsing sinusoidal and DC magnetic fields affect myosin phosphorylation in a cell-free preparation. Paper presented at the First World Congress on Electricity and Magnetism in Biology and Medicine.
- Martin AH (1988): Magnetic fields and time dependent effects on development. *Bioelectromagnetics* 9:393–396.
- Martin AH, Korall H, Forster B (1989): Magnetic field effects on activity and aging in honeybees. *J Comp Physiol* 164:423–431.
- Mastryukova VM, Rudneva SV (1978): Effect of a strong magnetostatic field on proliferation of duodenal epithelial cells in mice. *Biol Bull Acad Sci USSR* 5:371–374.
- Neurath PW (1968): High gradient magnetic field inhibits embryonic development of frogs. *Nature* 119:1358–1359.
- Neurath PW (1969): The effect of high-gradient, high-strength magnetic fields on the early development of frogs. In Barnothy MF (ed): “Biological Effects of Magnetic Fields, Vol. 2.” New York: Plenum Press.
- Nocente-McGrath C, McIsaac R, Ernst SG (1991): Altered cell fate in LiCl-treated sea urchin embryos. *Dev Biol* 147:445–450.
- Okazaki K (1975): Normal development to metamorphosis in the sea urchin embryo. In Czihak G (ed): “Biochemistry and Morphogenesis.” New York: Springer-Verlag.
- Pereira M, Nutini L, Fardon J, Cook E (1967): Effects of intermittent magnetic fields on cellular respiration. In Barnothy MF (ed): “Third International Biomagnetic Symposium.” Chicago: Chicago University Press.
- Pittman UG (1972): Biomagnetic responses in potatoes. *Can J Plant Sci* 52:727–733.
- Richardson BA, Yaga K, Reiter RJ, Morton DJ (1992): Pulsed static magnetic field effects on in-vitro pineal indoleamine metabolism. *Biochim Biophys Acta* 1137:59–64.
- Singh SS, Tiwari SP, Abraham J, et al. (1994): Magnetobiological effects on a cyanobacterium. *Electro Magnetobiol* 13:227–235.
- Tomlinson J, McGinty S, Kisin J (1981): Magnets curtail honey bee dancing. *Anim Behav* 29:307–308.
- Ueno S (1984): Embryonic development of frogs under strong DC magnetic fields. *IEEE Trans Magnet* 20:1663–1665.
- Verma SP, Goldner RB (1996): Raman spectroscopic evidence for structural changes in poly-L-lysine induced by an appropriate 50 mT static magnetic field. *Bioelectromagnetics* 17:33–36.
- Walker MM, Bitterman ME (1989): Conditioning analysis of magnetoreception in honeybees. *Bioelectromagnetics* 10:261–275.
- Weber T, Cerilli G (1971): Inhibition of tumor growth by the use of nonhomogeneous magnetic fields. *Cancer* 28:340–343.